

# Trauma Sensitivity Is Not Enough: Becoming a Trauma Competent Court

Brian L. Meyer, Ph.D.  
Interim Associate Chief, Mental Health Services/  
Supervisory Psychologist  
McGuire VA Medical Center  
Richmond, Virginia

October 29, 2015

# Disclaimer

The views expressed in this presentation are solely those of the presenter and do not represent those of the Veterans Administration, the Department of Defense, or the United States government.

**The author has no conflicts of interest to disclose.**

# What Is Trauma?

# What Do We Mean by “Trauma”?

Trauma results from an event, series of events, or set of circumstances that is experienced by an individual as physically or emotionally harmful or life threatening that has lasting adverse effects on the individual’s functioning and mental, social, emotional, or spiritual well-being.

SAMHSA, 2014

# What Do We Mean by “Trauma”?

Trauma results from an **event**, series of events, or set of circumstances that is experienced by an individual as physically or emotionally harmful or life threatening that has lasting adverse effects on the individual’s functioning and mental, social, emotional, or spiritual well-being.

SAMHSA, 2014

# What Do We Mean by “Trauma”?

Trauma results from an event, series of events, or set of circumstances that is **experienced** by an individual as physically or emotionally harmful or life threatening that has lasting adverse effects on the individual’s functioning and mental, social, emotional, or spiritual well-being.

SAMHSA, 2014

# What Do We Mean by “Trauma”?

Trauma results from an event, series of events, or set of circumstances that is experienced by an individual as physically or emotionally harmful or life threatening that has lasting adverse **effects** on the individual’s functioning and mental, social, emotional, or spiritual well-being.


SAMHSA, 2014

# Prevalence of Trauma in Justice-Involved Populations

The experience of trauma among people with substance abuse and mental health disorders, especially those involved with the justice system, is so high as to be considered *an almost universal experience*.

SAMHSA, 2013

# Maslow's Hierarchy of Needs




# Trauma and PTSD

# Incidence and Prevalence of Post-Traumatic Responses

- What is the relationship between trauma and PTSD?
- How common is PTSD?
- Which populations are more likely to develop PTSD?
- What are the symptoms of PTSD?
- Is all PTSD the same?
- What are the physical, psychological, moral, and social consequences of PTSD?

# Post-Traumatic Stress Disorder Is Characterized by:



- Exposure to a severe life-threatening event
- Repetitive re-experiencing of the LT event
- Avoidance of stimuli associated with trauma
- Negative mood and cognitions
- Increased arousal

# Post-Traumatic Responses Occur on a Continuum



None

Mild

Moderate

Severe

# The Continuum of Trauma

- When are you dealing with pre-existing behavior and when are you dealing with post-traumatic behavior?
  - When are you dealing with post-deployment behavior?
- What are the differences between acute, chronic, secondary, and complex trauma?
- Is it possible to have a post-traumatic response without developing PTSD?
- What is the range of post-traumatic responses?
- How do the type(s), frequency, and intensity of traumatic experiences affect an individual?
- What are the roles of environmental, historical and cultural traumas?

# The Long Reach of Childhood Experiences


You can spend a lifetime trying to forget a few minutes of your childhood.



HealthyPlace.com

- Are you familiar with the Adverse Childhood Experiences study?
- Do you use the ACE questionnaire?
- How do childhood traumatic experiences affect brain development?
- How do childhood traumatic experiences affect adult behavior?
- How do you know if this is happening in the defendant?

# THE SPECTRUM OF TRAUMA DISORDERS



Adjustment  
Disorder/  
PTS  
Symptoms

Sub-clinical  
PTSD

PTSD

Complex  
Trauma

Borderline and  
Antisocial  
Personality  
Disorders

Dissociative  
Identity  
Disorder

# Who Are We Kidding?




# The Importance of Cultural Context


- The culture can be traumatized
  - Historical trauma
  - Discrimination
- Trauma is understood through the lens of culture
  - What is the cultural meaning of traumatic events?
- What is the community's reaction to the trauma?
  - For example, returning military
  - For example, Hurricane Katrina

# **What You Need to Know: The Neurobiology of Trauma**

# Traumatic Stress and the Brain



# PTSD and the Brain



Amygdala – Emotional reactions, fight or flight alarm system

Hippocampus – Relay station for sorting memories


Prefrontal cortex – logic, reasoning, planning, impulse control, organizing

# For those that tell me,

*"PTSD? it's all in your head bro"...*



*Healthy Brain*



*PTSD*

## Yes, it is.

*Fight the phobia,  
End the STIGMA.*



[facebook.com/veteransptsdproject](https://facebook.com/veteransptsdproject)

Tom Kauffman '14

# The Effects of Abuse on Early Brain Development



Excess cortisol and heightened amygdalar response causing:

- Chronic fear and anxiety
- Inattention
- Overreactivity
- Impulsivity
- Hyperalertness and hyperarousal
- Sleep problems

# The Effects of Abuse on Early Brain Development

Increased epinephrine and stress steroids causing:

- Dissociation
- Disengagement
- Distorted attachments to others
- Numbing
- Emotional detachment
- Inability to feel empathy and remorse

Who does this sound like?

# PTSD: Increased Arousal

E. Trauma-related alterations in arousal and reactivity that began or worsened after the traumatic event:

- Irritable or aggressive behavior
- Self-destructive or reckless behavior
- Hypervigilance
- Exaggerated startle response
- Problems in concentration
- Sleep disturbance



# How Does the Neurobiology of Trauma Show Itself in the Courtroom?



- Agitation
- Anxiety and panic
- Hypervigilance
- Startle responses to noise
- Discomfort with crowds
- Being touched → alarm



- Distrust
- Defiance
- Disrespect
- Hostility
- Provocative

# **What You Need to Learn: Becoming Trauma Competent**

# What's the Real Story?



What they did to get into court



What happened to them to get them here

# Developmental Trauma

- What are Adverse Childhood Experiences (ACEs)?
- What are the effects of ACEs?
- What is the relationship between trauma and ACEs?
- How is developmental trauma different from adult trauma exposure?
- How are traumas passed from generation to generation?
- What is toxic stress and what are its effects?
- When does brain development end?

# Trauma

- What are the differences between interpersonal and environmental traumas?
- What are historical, intergenerational, and community traumas, and what are their effects?
- How are military traumas different than other traumas?
- What role does betrayal play in trauma?
- What are the differences between Big T and little t traumas?



# Trauma

## PEOPLE WHO HAVE EXPERIENCED TRAUMA ARE:



- What are the biological, psychological, spiritual, interpersonal, and societal effects of trauma?
- In what ways does the culture in which a person lives impact the experience of trauma?
- What is the relationship between trauma and emotional dysregulation?
- What conditions and problems are commonly co-morbid with trauma?

# Trauma and Substance Abuse

- In what ways are trauma and substance abuse related?
- How accurate is the self-medication hypothesis?
- How are substances used to cope to deal with the effects of trauma?



# Trauma Interventions

- What are trauma-responsive interventions and supports?
- What are the evidence-based assessments for trauma?
- Why don't current medications treat all of the symptoms of PTSD?
- Which current medications have been approved to treat some symptoms of PTSD?
- What are the evidence-based treatments for trauma?
- How do you know if a provider is actually using evidence-based treatments?
- Why is integrated treatment more successful than sequential or parallel treatment of trauma and substance abuse?

# Some Areas Where PTSD and the Legal System Intersect

- Domestic violence
- Child abuse
- Divorce
- Juvenile delinquency
- Homelessness
- DUI/DWI
- Misdemeanors
- Threats
- Violence
- Criminal actions



# **What You Need to Know: Adverse Childhood Experiences**

# The Adverse Childhood Experiences Study

- 17,421 adult patients of Kaiser Permanente
- Eight categories of adverse childhood events (ACEs) in the home: physical abuse, emotional abuse, sexual abuse, someone imprisoned, domestic violence, substance abuse, chronic mental illness, and loss of parent
- Results more than 50 years later:
  - More than 1/2 of population experienced one or more ACEs; 1/4 had two or more
  - The higher the ACE score, the worse the health and life outcomes

# Adverse Childhood Experiences

Felitti & D'Anda, 1998

ACEs = ADVERSE  
CHILDHOOD  
EXPERIENCES

*The three types of ACEs include*

ABUSE



Physical



Emotional



Sexual

NEGLECT



Physical



Emotional

HOUSEHOLD DYSFUNCTION



Mental Illness



Mother treated violently



Incarcerated Relative



Substance Abuse




Divorce

# ACE Study Results

- Greater health problems
- Greater behavioral health problems:
  - Alcoholism
  - Intravenous drug abuse
  - Smoking
  - Depression
  - Attempted suicide
- Greater occupational problems





**Mechanisms by Which Adverse Childhood Experiences Influence Health and Well-being Throughout the Lifespan**


# What You Need to Know: Complex Trauma

# What is Complex Trauma?

Complex psychological trauma results from exposure to severe stressors that (1) are repetitive or prolonged, (2) involve harm or abandonment by caregivers or other ostensibly responsible adults, and (3) occur at developmentally vulnerable times in the victim's life.

Ford and Courtois, 2009

# RESPONSE TO MULTIPLE TRAUMAS



After Bruce D. Perry, 1999


# Core Problems in Complex Trauma

- Affect dysregulation
- Dissociation
- Somatic dysregulation
- Impaired self-concept
- Disorganized attachment patterns

In addition to symptoms of PTSD and other comorbid disorders


Ford and Courtois, 2009

## What is Complex PTSD?




*Complex PTSD involves the core symptoms of PTSD plus additional groups of symptoms  
source: ISTSS Expert Consensus Treatment Guidelines For Complex PTSD In Adults*

**TraumaAndDissociation**  
[www.dissociative-identity-disorder.net/wiki/Complex\\_PTSD](http://www.dissociative-identity-disorder.net/wiki/Complex_PTSD)



# THE CATALYZING EFFECT OF COMPLEX TRAUMA



# **What You Need to Know: Common Co-Morbidities**

# Co-occurrence of PTSD and Substance Abuse


Co-occurring disorders are the rule rather than the exception.

(SAMHSA, 2002)



Figure 1

## COMORBID DISORDERS ARE THE RULE, RATHER THAN THE EXCEPTION FOR PTSD



(Chart: Kati Tal)

# Co-occurrence of PTSD and Substance Abuse

## National Comorbidity Survey

Among those with PTSD:

	Male	Female
Alcohol Abuse/ Dependence	51.9%	27.9%
Drug Abuse/ Dependence	34.5%	26.9%

Kessler et al., 1995

# Co-occurrence of PTSD and Substance Abuse

- PTSD and substance abuse co-occur at a high rate
  - 20-40% of people with PTSD also have SUDs (SAMHSA, 2007)
  - 40-60% of people with SUDs have PTSD
- Substance use disorders are 3-4 times more prevalent in people with PTSD than those without PTSD (Khantzian & Albanese, 2008)
- The presence of either disorder alone increases the risk for the development of the other
- PTSD increases the risk of substance relapse (Norman et al., 2007)
- The combination results in poorer treatment outcomes (Ouimette et al., 2003; Sonne et al., 2003)

# Co-Occurring PTSD and SUDs Make Each Other Worse

- Substance abuse exacerbates PTSD symptoms, including sleep disturbance, nightmares, rage, depression, avoidance, numbing of feelings, social isolation, irritability, hypervigilance, paranoia, and suicidal ideation
- People who drink or use drugs are at risk for being retraumatized through accidents, injuries, and sexual trauma

# Other Common Psychiatric Diagnoses in People Exposed to Traumatic Events

- Dysthymic Disorder
- Major Depressive Disorder
- Mood Disorder NOS
- Bipolar Disorder
- Generalized Anxiety Disorder
- Phobic Disorder
- Panic Disorder
- Schizophrenia
- Schizoaffective Disorder
- Anorexia
- Bulimia
- ADHD
- Oppositional Defiant Disorder
- Conduct Disorder
- Reactive Attachment Disorder
- Borderline Personality Disorder
- Antisocial Personality Disorder
- Narcissistic Personality Disorder

# Co-morbidity of PTSD and Pain

- **PTSD samples:**
  - 66-80% of individuals diagnosed with PTSD experience chronic pain (Beckham et al., 1997; Jakupcak, Osborne, Michael, Cook, Albrizio, & McFall, 2006; Shipherd et al., 2007)
- **Pain samples:**
  - 34% - 50% of civilians with chronic pain are diagnosed with PTSD (Geisser et al., 1996; Asmundson, et al., 1998)



# Insomnia and PTSD



- Insomnia is one of 20 defining characteristics of PTSD
  - It frequently continues even after PTSD is successfully treated
- Nightmares are another of the 20 defining characteristics of PTSD
  - Both nightmares and avoidance of nightmares can cause insomnia

# Being Trauma-Informed Is Not Enough

# What Does Being Trauma-Informed Mean?


- Being aware of the high frequency of trauma in defendants
  - 60% of people with substance abuse disorders have experienced trauma
  - The rate is probably much higher in judicial settings
- Rates of criminal behavior and violent offenses are much higher in victims of child abuse and neglect (Widom, 1989)
- Rates of child maltreatment are high among drug abusers
  - This is especially true among women, of whom 55-99% have a history of trauma (Najavits et al., 1997)
- Rape victims have far higher rates of drug abuse than those who have not been raped (Kilpatrick et al., 1992)
- 68% of prisoners report childhood abuse, and 23% report multiple forms of abuse (Weeks and Widom, 1998)

# What Does Being Trauma-Informed Mean?

- You understand that there is a link between trauma and substance abuse
- You also understand that, in order for substance abuse to end, trauma also needs to be treated
- Now you are trauma-informed.



# Trauma Informed Treatment ≠ Trauma Specific Treatment



- Trauma-informed treatment means that trauma is taken into account when treating substance abuse
  - *Beyond Trauma: A Healing Journey for Women* by Stephanie Covington
- Trauma specific treatment treats *both* trauma and substance abuse
- Trauma-specific treatment must be evidence-based
- Evidence-based means that research has shown treatment to be effective
  - *Seeking Safety* by Lisa Najavits

# **SAMHSA's Principles of Trauma Competency**

# The Four R's

- **Realize** the widespread impact of trauma and understand potential paths for recovery
- **Recognize** the signs and symptoms of trauma in participants, families, and staff
- **Respond** by integrating knowledge about trauma into policies, procedures, and practices
- Actively **resist re-traumatization**

# Principles of Trauma Competency

1. **Safety:** Staff, participants, and their families should feel physically and psychologically safe
2. **Trustworthiness and transparency:** Organizational operations and decisions are conducted with transparency and the goal of building and maintaining trust among staff, participants, and family members
3. **Peer support and mutual self-help:** Both are viewed as integral to the organizational and service delivery approach, and are understood as key vehicles for building trust, establishing safety, and empowerment


# Principles of Trauma Competency

4. **Collaboration and mutuality:** There is true partnering between staff and participants and among organizational staff from direct care to administrators
5. **Empowerment, voice, and choice:** In the organization and among staff, individual strengths are recognized, built on, and validated, and new skills are developed as necessary
6. **Cultural, historical, and gender issues:** The organization moves past cultural stereotypes and biases, and considers language and cultural considerations in providing support, offers gender-responsive services, leverages the healing value of traditional cultural and peer connections, and recognizes and addresses historical trauma

# Trauma Competency Means Changing Your Point of View

# The Traditional Approach to Criminal Justice


- The traditional approach can be re-traumatizing
  - Revolving door justice
  - Multigenerational justice
  - Disruption and violence in the courtroom
- How can we stop this cycle?



# What You See Depends on How You Look at It



# Continuum of Trauma Responsivity



# Central Tenets of Trauma Competency

---

1. Trauma is a public health problem

---

2. Assume that the defendant has experienced traumatic events

---

3. PTSD is a normal response to an abnormal event

---

4. Viewpoint changes from “What is wrong with you?” to “What happened to you?”

# Changing Your Approach

## Old View

- Trauma is irrelevant
- Trauma can be considered as a mitigating factor in sentencing
- See the problem behavior
- Respond to public pressure
- Needs of the institution

## New View

- Trauma is central
- Trauma-centric case processing
- See the whole person
- Respond to emerging science
- Needs of all participants

# Changing the Court's Approach

## Old Approach

- Adversarial
- Incarcerate
- Punishment
- Order
- Authoritarian

## New Approach

- Collaborative
- Treat
- Healing
- Partner
- Collaborative

# Changing Your Approach to Defendants

## Old Approach

- Tough love
- They are hopeless
- Judgmental
- Shames and blames
- Notices problems
- Defendant has a personality disorder
- Interprets behavior negatively

## New Approach

- Compassion
- We have hope
- Welcoming
- Accepts and holds accountable
- Notices strengths
- Defendant has experienced complex trauma
- Understands behavior is a communication and serves a function

# Changing Your Communication

## Hurtful

- Criticize
- Confront
- Sarcasm
- Talk loudly
- Distracted
- Judgmental
- Disrespectful
- Uses jargon

## Helpful

- Express concern
- Support
- Empathy
- Talk softly but firmly
- Active listening
- Accepting
- Patient
- Uses language everyone understands

# Changing Your Language

## Hurtful

- Characterizes behavior negatively, e.g., defendant is “disruptive and explosive”
- “You could stop using drugs if you wanted to.”
- “You should know better.”
- Victim

## Helpful


- Characterizes behavior constructively, e.g., defendant “needs calming strategies”
- “You need safety, stability, and support to succeed, and we want to help you.”
- “These are our expectations.”
- Survivor



**There is little or no cost to  
changing your approach.**

# Trauma Competency Means Changing Your Court

## Changing Your Point of View: 5 Ps, an E, and an A



Becoming a trauma competent court requires major shifts in your environment, philosophy, attitudes, perspective, policies, procedures, and practices

# Reconstruct the Physical Environment

- The goal is to reduce environmental stress
- Build buildings with easy navigation
- Smaller rooms are better
- Everyone sits at the same table
  - The judge joins
- Have separate waiting rooms for alleged perpetrator and trauma survivor





# Reconstruct the Environment

- Avoid ticking clocks and loud noises
- No yelling
- Keep the temperature comfortable



# Decrease Perceived Threats



- Bailiffs should not stand behind defendants
- Respect personal space
  - No touching
- Avoid trauma triggers *when possible*
  - No handcuffs or shackles
  - Avoid jumpsuits
  - Don't put defendants in isolation rooms

# Take Steps to Avoid Re-traumatization of Participants

- Decrease the power dynamic
  - Judge comes down from the bench
  - Judge takes off robe
- Use a solution-oriented approach instead
  - “What can you do differently? How can other people help?”
- Create a solution-oriented team
  - Invite everyone to participate actively
  - This is empowering




# Trauma in the Courtroom: What You Can Do

# What You Look Like to Them



# You Have to Change Their Perceptions, Too



Mehrabian & Ferris, 1967

- What are your facial expressions?
- What are you communicating non-verbally?
- What is your tone of voice?
- What is the volume of your voice?
- How do you respond to their behavior?

# Four Things You Must Establish Above All

Safety

Trust

Respect

A sense of some control

# **What You Need to Know: Evidence-Based Assessment and Treatment**


# Evidence-Based Assessments for Trauma

- PTSD Checklist 5 (PCL 5)
  - 20 item checklist corresponding to 20 symptoms of PTSD in DSM 5
- Clinician Assessment of PTSD Symptoms (CAPS)
  - This is the gold standard of PTSD assessment

# Medical Treatment of Trauma

- Medication for symptom management and co-morbid disorders
  - Antidepressants
  - Mood stabilizers
  - Anticonvulsants
  - Sleep aids, including Prazosin for nightmares
  - Atypical antipsychotics **No longer**
  - Anxiolytics **Not benzodiazepines**
- Only SSRIs are approved for treating PTSD
- There is no medication that specifically “cures” PTSD

# Phases of Integrated Treatment



After Herman, 1992

# Stage I: Safety and Stabilization

- Alliance building
- Psychoeducation about multiple traumas
- Safety
- Stabilization
- Skills-building
  - Affective regulation
  - Cognitive
  - Interpersonal
- Self-care



# Stage I: Safety

- Safety plans
- Tension reduction activities (e.g., exercise)
- Harm reduction and elimination
  - Self-harm and suicidal behaviors
  - Gambling
  - Driving
  - Fighting
  - Eating
  - Sex
  - Medication
  - Breaking laws



# Harm Reduction for Co-occurring Trauma and Substance Abuse in Drug Courts

- *Immediate* abstinence may not be possible for complex trauma patients
  - Alcohol and drugs are frequently used to regulate emotions
  - If they are taken away completely, they may be flooded by traumatic memories and emotions, which may lead to relapse
- Gradual reduction of substances with a goal of abstinence may be more realistic
- Punishing them for harm reduction is countertherapeutic and may result in failure

# Stage I: Stabilization

- Elimination of drug and alcohol abuse
- Health
- Housing
  - In a safe neighborhood
- Income
  - Employment
  - Financial skills (budgeting, banking)
- Transportation
- Setting and keeping a schedule



# Requiring AA/NA May Fail for People with Complex Trauma


- The first step in AA/NA is to admit helplessness
  - This may reignite their traumas
- AA/NA requires acknowledgement of a higher power
  - People who have experienced complex trauma may be agnostic or atheistic
- Therefore, other groups like SMART Recovery may fit better

# Evidence-Based Treatments for Stage I

- Seeking Safety
- Dialectical Behavior Therapy (DBT)
- Mindfulness-Based Stress Reduction
- Therapies for specific problems
  - Imagery Rehearsal Therapy
  - Cognitive-Behavioral Therapy
  - Motivational Interviewing
  - SAMHSA's Anger Management workbook

# Seeking Safety

- An integrated treatment for PTSD and Substance Abuse
- Combines psychoeducational and psychodynamic treatment
- 25 lessons on topics that overlap between PTSD and Substance Abuse
  - Safety Skills
  - Grounding
  - Anger
  - Boundaries
  - Self-care
  - Honesty
  - Compassion




# Seeking Safety

- Can be provided by professionals or paraprofessionals
- 6 randomized controlled trials and 3 controlled studies
- Seeking Safety has shown positive results across all studies (Najavits & Hien, 2013)
- Seeking Safety is the only evidence-based treatment for PTSD and Substance Abuse



# DBT Skills Training

- Four topics with multiple lessons
  - Mindfulness
  - Interpersonal Effectiveness
  - Distress Tolerance
  - Affect Regulation
- New manual provides suggested menus of different specific skills and exercises with different populations




# DBT Results


- 18 randomized controlled trials
- Results are all positive
- Populations include:
  - Women: with Borderline Personality Disorder (BPD) and suicidality, with BPD and substance dependence, with bulimia nervosa, with binge eating disorder, with opiate-addiction and BPD, domestic violence victims, with childhood sexual abuse, and with trichotillomania;
  - Adults: with BPD, with personality disorders, with Bipolar Disorder, prisoners with intellectual disabilities, and prisoners with impulsivity;
  - Male prisoners; and
  - Adolescents: suicidal, female offenders, with self-injurious behavior, with eating disorders

# CBT-I for Insomnia

- Cognitive-Behavioral Therapy for Insomnia (Perlis et al., 2008)
  - Psychoeducation about sleep and what interferes with it
  - Sleep restriction
  - Stress management
  - Cognitive restructuring
  - Relapse prevention



# Evidence-Based Stage II Treatments for PTSD



# Cognitive Processing Therapy

- A cognitive intervention to change the way a traumatized person thinks
- 12 weekly sessions delivered in a structured, manualized protocol
  - Number of sessions can be expanded
- May or may not include a trauma narrative
- Can be delivered individually and/or in groups
- Homework worksheets between sessions

# Cognitive Processing Therapy

- Central techniques:
  - Identifies stuck points
  - Examines evidence for thoughts and beliefs
  - Challenges beliefs
- Changing the interpretation of the traumatic event changes the emotions resulting from the event
- CPT is an effective treatment for PTSD (Vickerman & Margolin, 2009; Ougrin, 2011; Jonas et al., 2013; Ehring et al., 2014)
- CPT successfully treats complex trauma (Resick et al., 2003; Galovski et al., 2013)

# Prolonged Exposure

- A behavioral intervention that repeatedly exposes patients to distressing stimuli in order to decrease their anxiety in response to those stimuli
- 10 weekly sessions
- First part involves *in vivo* exposure to places that increase anxiety (e.g., public places)
  - Uses an anxiety hierarchy

# Prolonged Exposure

- Second part involves writing and dictating a trauma narrative focusing on one traumatic experience
  - The patient listens to the narrative over and over for an hour each day
  - Repeated and prolonged exposure decreases their anxiety
- ◆ Prolonged exposure is an effective treatment for PTSD (Vickerman & Margolin, 2009; Ougrin, 2011; Jonas et al., 2013; Ehring et al., 2014)
- There is no evidence that it successfully treats complex trauma
- The evidence shows that it does not effectively treat substance abuse, even when a substance abuse program is provided side by side

# Eye Movement Desensitization and Reprocessing

- Patient focuses on distressing image
  - States a belief that goes with it
  - Notices feelings that go with it
  - Identifies body sensations that go with it
- Therapist passes fingers back and forth, guiding the eyes
- As this occurs, the images, thoughts, feelings, and body sensations change
- Adaptive information processing results

# Eye Movement Desensitization and Reprocessing

- Auditory and tactile alternatives to eye movements using bilateral stimulation
- Additional exercises:
  - Safe Place
  - Lightstream
  - Resource-building
  - Protocol for substance abuse
  - Etc.



# Eye Movement Desensitization and Reprocessing Results

- EMDR works for PTSD and complex trauma (Davidson & Parker, 2001; Foa et al., 2009; Maxfield & Hyer, 2002; Seidler & Wagner, 2006)
- EMDR addresses substance abuse (Vogelmann-Sine et al., 1998)
- EMDR uses the same mechanism for resolution (eye movements) that sleep does
  - EMDR also targets nightmares

# Promising Treatments: STAIR Narrative Therapy

Skills Training in Affective and Interpersonal Regulation (STAIR)  
Narrative Therapy (Cloitre et al., 2006)

- Uses coping skills from Stress Inoculation Training and Dialectical Behavior Therapy
- 8-10 sessions of skills building and 8 sessions of narrative therapy
- This is the only Phase I and Phase II treatment for complex trauma

# Promising Treatments: STAIR Narrative Therapy

- Narrative Therapy: developing an autobiography
  - Repeated narration to organize trauma memory and reduce fear
  - Analyze meaning of event(s) to revise beliefs/schemas about self and others, integrate traumatic memories into a life history, and explore and resolve feelings other than fear
  - Continue practice of STAIR skills

# Promising Treatments: STAIR Narrative Therapy

- Four studies of STAIR Narrative Therapy (Cloitre et al., 2002; Levitt et al., 2007; Trappler & Newville, 2007; Cloitre et al., 2010) show:
  - Decreases in PTSD symptoms
  - Improvements in interpersonal problems
  - Improvements in emotion regulation
- Studies of women with child abuse histories, post 9/11 survivors, and inpatients with co-morbid PTSD and Schizoaffective Disorder

## Stage III: Reconnection



- Gradually decrease isolation
- Re-establishing estranged relationships
- Developing trusting relationships
- Developing intimacy
- Developing sexual intimacy
- Parenting
- Community-based activities
- Spirituality

## Stage III: Reconnection

- Giving back to the community
- Making amends
- Acceptance
- Reclaiming
- Creativity
- Finding meaning
- Post-traumatic growth

**“IT’S NOT TOO LATE TO DEVELOP NEW  
FRIENDSHIPS OR RECONNECT WITH PEOPLE.”**

**MORRIE SCHWARTZ**

© Lifehack Quotes

# Wellness Activities

- Mindfulness Meditation
- Yoga
- Qi Gong
- Tai Chi
- Massage
- Acupuncture



# Trauma in the Courtroom: Secondary Traumatization

# Secondary Traumatization

- Secondary traumatization typically occurs when a person hears stories of traumatic experiences
- It may also occur when a person lives with someone who has been traumatized
- It results in post-traumatic symptoms like hypervigilance, irritability, startle responses, distrust of others, negative thoughts/feelings/ beliefs, avoidance, and isolation
- Children are particularly susceptible to Secondary Traumatization

# Dealing with Secondary Traumatization

- Ongoing training
- Regular scheduled debriefing meetings
- Use of EAP or psychotherapy
- Set firm boundaries between work and home
- Engage your support network
- Play
- Be creative: sing, dance, write, draw, sculpt, etc.

# Resources

# Trauma Competent Courts

- *Essential Components of Trauma-Informed Judicial Practice*, SAMHSA. Retrieved from [http://www.nasmhpd.org/sites/default/files/JudgesEssential\\_5%201%202013finaldraft.pdf](http://www.nasmhpd.org/sites/default/files/JudgesEssential_5%201%202013finaldraft.pdf)
- Also valuable: *TIP 57: Trauma-Informed Care in Behavioral Health Services*, SAMHSA, available at [www.store.samhsa.gov](http://www.store.samhsa.gov).
- SAMHSA [www.samhsa.gov](http://www.samhsa.gov)

# Adverse Childhood Experiences Study

- [www.cdc.gov/ace](http://www.cdc.gov/ace)
- <http://acestudy.org>
- ACE questionnaire  
[http://acestudy.org/yahoo\\_site\\_admin/assets/docs/ACE\\_Calculator-English.127143712.pdf](http://acestudy.org/yahoo_site_admin/assets/docs/ACE_Calculator-English.127143712.pdf)

## Resources for PTSD

- *Handbook of PTSD* by Matthew Friedman, Terence Keane, and Patricia Resick
- *Once a Warrior, Always a Warrior: Navigating the Transition from Combat to Home--Including Combat Stress, PTSD, and mTBI* by Charles Hoge
- *When Someone You Love Suffers from Posttraumatic Stress: What to Expect and What You Can Do* by Claudia Zayfert and Jason Deviva

# Resources for PTSD

- National Center for PTSD: [www.ptsd.va.gov](http://www.ptsd.va.gov)
- International Society for Traumatic Stress Studies: [www.istss.org](http://www.istss.org)
- International Society for the Study of Trauma and Dissociation:  
[www.isst-d.org](http://www.isst-d.org)
- PTSD 101 courses:  
[www.ptsd.va.gov/professional/ptsd101/course-modules.asp](http://www.ptsd.va.gov/professional/ptsd101/course-modules.asp)

# Resources for Complex Trauma

- Trauma and Recovery, 1992, Judith Herman
- Luxenberg, T., Spinazzola, J., and van der Kolk, B. (2005). Complex Trauma and Disorders of Extreme Stress (DESNOS) Diagnosis, Part One: Assessment (2005). Directions in Psychiatry, 21, 373-393.
- Treating Complex Traumatic Stress Disorders, 2009, Christine Courtois and Julian Ford, eds.
- Treatment of Complex Trauma: A Sequenced, Relationship-Based Approach (2012), Christine Courtois, Julian Ford, and John Briere
- <http://www.nctsn.org/trauma-types/complex-trauma/assessment>

# Resources

- *Complex Trauma in Children and Adolescents*, NCTSN White Paper, available at [http://www.nctsn.org/sites/default/files/assets/pdfs/Complex Trauma All.pdf](http://www.nctsn.org/sites/default/files/assets/pdfs/Complex_Trauma_All.pdf)
- [The Trauma Recovery Group: A Guide for Practitioners \(2011\)](#), Michaela Mendelsohn, Judith Herman, Emily Schatzow, and Diya Kallivayalil
- International Society for Traumatic Stress Studies: <http://www.istss.org>
- Trauma Focused-Cognitive Behavioral Therapy: <http://tfcbt.musc.edu>

# PTSD and SUDs

- PTSD 101 course about treating PTSD and SUDs: [www.ptsd.va.gov/professional/ptsd101/course-modules/SUD.asp](http://www.ptsd.va.gov/professional/ptsd101/course-modules/SUD.asp)
- Practice recommendations for treating co-occurring PTSD and SUDs: [www.ptsd.va.gov/professional/pages/handouts-pdf/SUD\\_PTSD\\_Practice\\_Recommend.pdf](http://www.ptsd.va.gov/professional/pages/handouts-pdf/SUD_PTSD_Practice_Recommend.pdf)

# Resources for PTSD and SUDS

- *Trauma and Substance Abuse (2<sup>nd</sup> ed.)* by Page Ouimette and Jennifer Read
- *Treating Survivors of Childhood Abuse: Psychotherapy for the Interrupted Life* by Marylene Cloitre, Lisa Cohen, and Karestan Koenen
- *Concurrent Treatment of PTSD and Substance Use Disorders Using Prolonged Exposure (COPE) Therapist Guide* by Sudie Back, Edna Foa, Therese Killeen, Katherine Mills, Maree Teesson, Bonnie Cotton, Kathleen Carroll, and Kathleen Brady

# Seeking Safety

- Seeking Safety (1998), Lisa Najavits
- *8 Keys to Trauma and Addiction Recovery* (2015), Lisa Najavits
- <http://www.treatment-innovations.org/seeking-safety.html>

# Dialectical Behavior Therapy

- Cognitive-Behavioral Treatment of Borderline Personality Disorder (1993), Marsha Linehan
- DBT Skills Training Manual, 2<sup>nd</sup> Ed. (2014), Marsha Linehan
- DBT Skills Training Handouts and Worksheets, 2<sup>nd</sup> Ed. (2014), Marsha Linehan
- <http://www.behavioraltech.com>
- <http://www.linehaninstitute.org/>

# Prolonged Exposure

- Prolonged Exposure Therapy for PTSD: Emotional Processing of Traumatic Experiences Therapist Guide (2007), Edna Foa, Elizabeth Hembree and Barbara Olaslov Rothbaum
- Reclaiming Your Life from a Traumatic Experience: A Prolonged Exposure Treatment Program Workbook (2007), Barbara Rothbaum, Edna Foa and Elizabeth Hembree

# Cognitive Processing Therapy

- Cognitive Processing Therapy for Rape Victims: A Treatment Manual (1993), Patricia Resick and Monica Schnicke

# EMDR

- Eye Movement Desensitization and Reprocessing (EMDR): Basic Principles, Protocols, and Procedures, 2<sup>nd</sup> Ed. (2001), Francine Shapiro
- Getting Past Your Past: Take Control of Your Life with Self-Help Techniques from EMDR Therapy (2013), Francine Shapiro
- [www.emdr.com](http://www.emdr.com)
- [www.emdria.org](http://www.emdria.org)
- [www.emdrhap.org](http://www.emdrhap.org)

# STAIR Narrative Therapy



- Treating Survivors of Childhood Abuse: Psychotherapy for the Interrupted Life (2006), Marilene Cloitre, Lisa Cohen, and Karestan Coenen
- Online at <http://www.stairnt.com/index.html>  
[http://www.ptsd.va.gov/professional/continuing\\_ed/STAIR\\_online\\_training.asp](http://www.ptsd.va.gov/professional/continuing_ed/STAIR_online_training.asp)

# Family Resources

- *When Someone You Love Suffers from Posttraumatic Stress: What to Expect and What You Can Do* by Claudia Zayfert and Jason Deviva
- *Finding My Way: A Teen's Guide to Living with a Parent Who Has Experienced Trauma* (2005), Michelle Sherman and DeAnne Sherma
- <http://www.ptsd.va.gov/public/pages/fslist-family-relationships.asp>

# SMART Recovery

- [www.smartrecovery.org](http://www.smartrecovery.org)
- <http://smartrecoverytraining.org/moodle/>
- <http://www.smartrecovery.org/community/#.Vims8GtRI2Y>



SMART  
Recovery App



# Cognitive-Behavioral Therapy for Insomnia

- *Cognitive Behavioral Treatment of Insomnia: A Session-by-Session Guide* (2008), by Michael L. Perlis, Carla Jungquist, Michael Smith, and Donn Posner
- *Overcoming Insomnia: A Cognitive-Behavioral Therapy Approach Workbook* (2008), by Jack Edinger and Colleen Carney

# Online Resources

- Self-assessment Mental Health screening

<http://www.militarymentalhealth.org/>

- Problem-solving

<http://startmovingforward.t2.health.mil/>

- Wellness resources

<http://afterdeployment.t2.health.mil/>

# Online and Telephone Resources




**Veterans  
Crisis Line**  
1-800-273-8255 PRESS 1

**IT'S YOUR CALL**


Confidential help for  
Veterans and their families

• • • • Confidential chat at [VeteransCrisisLine.net](http://VeteransCrisisLine.net) or text to 838255 • • • •



LEARN FROM  
VETERANS HOW  
PTSD TREATMENT  
CAN TURN YOUR  
LIFE AROUND

**ABOUT FACE**



**MAKE THE CONNECTION**  
Share your experiences and support for Veterans.

Home | Who You Are | Life Events & Experiences | Video Gallery | Signs & Symptoms | Conditions | Resources & Support

WELCOME TO  
MAKE THE CONNECTION

Connecting Veterans and their friends and family members with information, resources, and solutions to issues after they leave service.

**CUSTOMIZE THIS SITE FOR YOU**

I am  MALE  FEMALE

I served during:

I served in:


I was exposed to combat  YES  NO

**CUSTOMIZE NOW**

# Self-Help Mobile Applications

<http://www.t2health.org/mobile-apps>

- PTSD Coach



- T2 MoodTracker



- Breathe 2 Relax



- Tactical Breather



- LifeArmor (includes family section)




# Self-Help Mobile Applications

- Positive Activity Jackpot



<http://www.militarymentalhealth.org/articles/media/>

- Virtual Hope Box
- Provider Resilience
- More to come!




# Mobile Applications That Assist Psychotherapy

- PE Coach



- CPT Coach



- CBT-I Coach



- Mindfulness Coach



- ACT Coach



**Contact:**  
**Brian L. Meyer, Ph.D.**  
[Brian.Meyer@va.gov](mailto:Brian.Meyer@va.gov)